FORM PTO-1449 (Modified) INFORMATION DISCLOSURE STATEMENT BY APPLICANT

U.S. Department of Commerce Patent and Trademark Office

Attorney Docket No.: STRATA-06948

Serial No.: 10/087,388

INFORMATION DISCLOSURE STATEMENT BY APPLICANT (37 CFR § 1.98(b))			Applicant: Allen Comer et al.	
			Filing Date: 03/01/02	Group Art Unit:
	r	OTHER DOCUMENTS (Including Author, Title, I	Date, Relevant Pages, Place of Pu	ublication)
	1	Berger et al., Secreted placental alkaline phosphatase: a pow 66:1-10 (1998)	verful new quantitative indicator of	of gene expression in eukaryotic cells, Gene
	2	Jaspers et al., Arsenite Exposure of Cultured Airway Epithelial Cells Activates κB-dependent Interleukin-8 Gene Expression in the Absen of Nuclear Factor κB Nuclear Translocation, J. Biol. Chem. 274:31025-31033 (1999)		
	3	Muller et al., Enhanced expression of IL-8 in normal human keratinocytes and human keratinocyte cell line HaCaT in vitro after stimulativith contact sensitizers, tolerogens and irritants, Exp. Dermatol. 3:298-303 (1994)		
•	4	Newby et al., Cytokine Release and Cytotoxicity in Human Keratinocytes and Fibroblasts Induced by Phenols and Sodium Dodecyl Sulfa Soc. of Investigative Dermatology, 115:292-298 (2000)		
	5	Fisher et al., All-Trans Retinoic Acid Induces Cellular Retinol-Binding Protein in Human Skin In Vivo, Soc. of Investigative Dermatology 105:80-86 (1995)		
	6	Wilmer et al., Cytokine Induction in Human Epidermal Keratinocytes Exposed to Contact Irritants and Its Relation to Chemical-Induced Inflammation in Mouse Skin, Soc. of Investigative Dermatology, 102:915-922 (1994)		
	7 Roguet, Cell Biology and Toxicology, 15:64-75 (1999)			
	8	van Ruissen et al., Differential Effects of Detergents on Keratinocyte Gene Expression, Soc. of Investigative Dermatology, 110:358-363 (1998)		
	9	Terunuma et al., Cytokine mRNA profiles in cultured human epicutaneous stimuli induced by skin barrier perturbation in c Science, 26:85-93 (2001)		
	10	Medina et al., Use of Human Skin Equivalent Apligraf for in Toxicology and Applied Pharmacology, 164:38-45 (200)	Vitro Assessment of Cumulative	Skin Irritation Potential of Topical Product
	11	Muller-Decker et al., Keratinocyte-Derived Proinflammatory Possible Alternative to the Draize Skin Irritation Test, Toxico	key Mediators and Cell Viability blogy and Applied Pharmacology,	as in Vitro Parameters of Irritancy: A, 127:99-109 (1994)
	12	Abe et al., Interleukin-8 Gene Repression by Clarithromycin Is Mediated By The activator Protein-1 Binding Site in Human Bronchial Epithelial Cells, Am. J. Respir. Cell Mol. Biol., 22:51-60 (2000)		
			· · · · · · · · · · · · · · · · · · ·	
aminer:		/Leon Lankford Jr/	Date Considered:	01/21/2009